

Literature CA72-4

- 1: Ohuchi N, Takahashi K, Matoba N, Mori S. [A radioimmunoassay for circulating tumor-associated glycoprotein (TAG-72) using the antigen determinant CA 72-4]. *Gan To Kagaku Ryoho*. 1988 Sep;15(9):2761-6.
- 2: Ohuchi N, Matoba N, Taira Y, Takahashi K, Sakai N, Sato K, Fujita N, Mochizuki F, Nishihira T, Mori S. [Levels of circulating tumor-associated glycoprotein (TAG-72) in patients with carcinoma using a novel tumor marker, CA 72-4]. *Gan To Kagaku Ryoho*. 1988 Sep;15(9):2767-72.
- 3: Ferroni P, Szpak C, Greiner JW, Simpson JF, Guadagni F, Johnston WW, Colcher D. CA 72-4 radioimmunoassay in the diagnosis of malignant effusions. Comparison of various tumor markers. *Int J Cancer*. 1990 Sep 15;46(3):445-51.
- 4: Guadagni F, Roselli M, Cosimelli M, Spila A, Cavaliere F, Tedesco M, Arcuri R, Abbolito MR, Casale V, Pericoli MN, Vecchione A, Casciani CU, Greiner JW, Schlom J. Correlation between tumor-associated glycoprotein 72 mucin levels in tumor and serum of colorectal patients as measured by the quantitative CA 72-4 immunoassay. *Cancer Res*. 1996 Nov 5;56(22):5293-8.
- 5: Gadducci A, Ferdeghini M, Ceccarini T, Prontera C, Facchini V, Bianchi R, Fioretti P. The serum concentrations of TAG-72 antigen measured with CA 72-4 IRMA in patients with ovarian carcinoma. Preliminary data. *J Nucl Med Allied Sci*. 1989 Jan-Mar;33(1):32-6.
- 6: Reinsberg J, Gast B, Schmolling J, Wagner U, Krebs D. Specific interference with the determination of the tumour-associated glycoprotein 72 by human anti-idiotypic antibodies formed after treatment with the anti-tumour-associated glycoprotein 72 antibody B72.3. *Eur J Clin Chem Clin Biochem*. 1994 Sep;32(9):691-6.
- 7: Jacobs IJ, Rivera H, Oram DH, Bast RC Jr. Differential diagnosis of ovarian cancer with tumour markers CA 125, CA 15-3 and TAG 72.3. *Br J Obstet Gynaecol*. 1993;100(12):1120-4.
- 8: Contegiacomo A, Alimandi M, Muraro R, Pizzi C, Calderopoli R, De Marchis L, Sgambato A, Pettinato G, Petrella G, De Filippo MR, et al. Expression of epitopes of the tumour-associated glycoprotein 72 and clinicopathological correlations in mammary carcinomas. *Eur J Cancer*. 1994;30A(6):813-20.
- 9: Filella X, Molina R, Jo J, Bedini JL, Joseph J, Ballesta AM. Tumor associated glycoprotein-72 (TAG-72) levels in patients with non-malignant and malignant disease. *Bull Cancer*. 1992;79(3):271-7. Review.
- 10: Kuroki M, Fernsten PD, Wunderlich D, Colcher D, Simpson JF, Poole DJ, Schlom J. Serological mapping of the TAG-72 tumor-associated antigen using 19 distinct monoclonal antibodies. *Cancer Res*. 1990 Aug 15;50(16):4872-9.

11: Motoo Y, Satomura Y, Kawakami H, Watanabe H, Ohta H, Okai T, Sawabu N. Serum levels of tumor-associated glycoprotein (TAG-72) in digestive cancers. *Oncology* 1990;47(6):456-62.

12: Gero EJ, Colcher D, Ferroni P, Melsheimer R, Giani S, Schlom J, Kaplan P. CA 72-4 radioimmunoassay for the detection of the TAG-72 carcinoma-associated antigen in serum of patients. *J Clin Lab Anal.* 1989;3(6):360-9.

13: Sheer DG, Schlom J, Cooper HL. Purification and composition of the human tumor-associated glycoprotein (TAG-72) defined by monoclonal antibodies CC49 and B72.3. *Cancer Res.* 1988 Dec 1;48(23):6811-8.

14: Muraro R, Kuroki M, Wunderlich D, Poole DJ, Colcher D, Thor A, Greiner JW, Simpson JF, Molinolo A, Noguchi P, et al. Generation and characterization of B72.3 second generation monoclonal antibodies reactive with the tumor-associated glycoprotein 72 antigen. *Cancer Res.* 1988 Aug 15;48(16):4588-96.

15: Johnson VG, Schlom J, Paterson AJ, Bennett J, Magnani JL, Colcher D. Analysis of a human tumor-associated glycoprotein (TAG-72) identified by monoclonal antibody B72.3. *Cancer Res.* 1986 Feb;46(2):850-7.